

Article PDF Available

نقش ویژگیهای شخصیتی در پیش‌بینی اختلال استرس پس از سانحه در افراد در معرض آسیب

August 2018

Project: [Treatment of mathematical disorders](#)**Authors:****Mohammad Narimani**

University of Mohaghegh Ardabili

**Sajjad Basharpour**

University of Mohaghegh Ardabili

[Download full-text PDF](#)[Read full-text](#)[Download citation](#)[Copy link](#)[References \(23\)](#)**Abstract**

This research has been carried out to study the role of personality traits in predicting posttraumatic stress disorder (PTSD) in people who are exposed to psychological trauma. The total number of ER nurses and fire department staff members in Urumia city constituted the statistical population for of this research. From each group 50 individuals were randomly selected, yielding a sample of 100 participants for the research. Participants were tested individually by administering the Mississippi Posttraumatic Stress Disorder Scale and NEO personality inventory. The collected data were analyzed by MANOVA and regression tests. Results showed a significant difference in neuroticism, openness to experience, and conscientiousness of the two groups. However, there were no significant differences in

Discover the world's research

- 20+ million members
- 135+ million publications
- 700k+ registered projects

[Join for free](#)[Public Full-text \(1\)](#)

Content uploaded by [Mohammad Narimani](#) Author content
Content may be subject to copyright.

Psychological Research, 2010
Vol. 13, No. 1

پژوهشنامه‌ای روان‌شناسی، ۱۳۸۹،
دوره ۱۳، شماره ۱

نقش ویژگی‌های شخصیتی در پیش‌بینی اختلال استرس پس از سانحه در افراد در معرض آسیب

Role of Personality Traits in the Prediction of Posttraumatic Stress Disorder in People Exposed to Trauma

Mohammad Narimani, Ph.D.*

University of Mohaghegh Ardebili

Sajjad Basharpoor, M.A.

University of Mohaghegh Ardebili

دکتر محمد نریمانی

دانشگاه محقق اردبیلی

سجاد بشربور

دانشگاه محقق اردبیلی

Abstract: This research has been carried out to study the role of personality traits in predicting posttraumatic stress disorder (PTSD) in people who are exposed to psychological trauma. The total number of ER nurses and fire department staff members in Urumia city constituted the statistical population for of this research. From each group 50 individuals were randomly selected, yielding a sample of 100 participants for the research. Participants were tested individually by administering the Mississippi Posttraumatic Stress Disorder Scale and NEO personality inventory. The collected data were analyzed by MANOVA and regression tests. Results showed a significant difference in neuroticism, openness to experience, and conscientiousness of the two groups. However, there were no significant differences in

چکیده: تحقیق حاضر با هدف بررسی نقش ویژگی‌های شخصیتی در پیش‌بینی اختلال استرس پس از سانحه در افرادی که در معرض تروما هستند انجام گرفت. کلیه برسitaran اورژانس بیمارستان‌ها و کارکنان مرکز آتش‌نشانی شهرستان ارومیه جامعه آماری این پژوهش را تشکیل می‌دادند. از هر گروه تعداد ۵۰ نفر به روش نمونه‌گیری تصادفی ساده انتخاب شدند و در کل ۱۰۰ نفر حجم نمونه این پژوهش را تشکیل دادند. کلیه آزمودنی‌ها به پرسشنامه شخصیتی NEO و مقیاس اختلال استرس پس از سانحه می‌سینی پاسخ دادند. داده‌های به دست آمده با استفاده از آزمون‌های تحلیل واریانس چندمتغیری و تحلیل رگرسیون چندگانه مورد تجزیه و تحلیل قرار گرفتند. نتایج حاصل نشان داد که بین افراد در معرض تروماتی مبتلا به اختلال استرس پس از سانحه و افراد در معرض تروماتی سالم، در ویژگی‌های شخصیتی روان‌نگری، گشودگی به تجربه و

* narimani@uma.ac.ir

دریافت اصلاح نهایی مقاله: ۱۳۸۹/۲/۵

پذیرش مقاله: ۱۳۸۹/۳/۱۹

[Citations \(0\)](#)[References \(23\)](#)

Post-traumatic stress and associated disorders among Vietnam veterans: The significance of combat exposure and social support

[Full-text available](#) Article

Apr 1995

Joseph A Boscarino

[View](#) [Show abstract](#)

Traumatic events and post-traumatic stress disorder in an urban population of young adults

Article

Jan 1991

B. Breslau · G. Davis · P Andreski · E. Peterson

[View](#)

Prewar factors in combat-related posttraumatic stress disorder: Structural equation modeling with a national sample of female and male Vietnam veterans

Article

Jun 1996

Daniel W King · L.A. King · David W. Foy · David M. Gudanowski

[View](#) [Show abstract](#)

Does personality moderate the relationship between stress and health behavior? Expanding the nomological network of the five-factor model

Article

Dec 2008

Dave Korotkov

[View](#) [Show abstract](#)

Personality and PTSD II: Personality assessment of PTSD-diagnosed Vietnam veterans using the Cloninger Tridimensional Personality Questionnaire (TPQ)

Article

Jan 1997 · DEPRESS ANXIETY

Harvey Richman · B Christopher Frueh

[View](#) [Show abstract](#)

The relationship between neuroticism, pre-traumatic stress, and post-traumatic stress: A prospective study

Article

Jul 2003 · PERS INDIV DIFFER

Iris M. Engelhard · Marcel van den Hout · Merel Kindt

[View](#) [Show abstract](#)

The Aetiology of Post-traumatic Morbidity: Predisposing, Precipitating and Perpetuating Factors

Article

Mar 1989

Alexander Mcfarlane

[View](#) [Show abstract](#)

NEO-PI profiles in PTSD as a function of trauma level

[Article](#)

Oct 1993

F. Suzanne Talbert · Lorraine C. Braswell · J. William Albrecht · Patrick A. Boudewyns

[View](#) [Show abstract](#)**Personality Disorders in Treatment-Seeking Combat Veterans with Posttraumatic Stress Disorder**[Article](#)

Aug 1993

S M Southwick ·  Rachel Yehuda · Earl L. Giller[View](#) [Show abstract](#)**Childhood physical abuse and combat-related posttraumatic stress disorder in Vietnam veterans**[Article](#)

Mar 1993

 James Douglas Bremner · S M Southwick · D R Johnson ·  Dennis Charney[View](#) [Show abstract](#)[Show more](#)

Recommendations [Discover more](#)**Project**

Treatment of mathematical disorders

Malahat Amani · Mohammad Narimani

[View project](#)**Project**

samadi

Deleted Profile · Seyfollah Aghajani · Mohammad Narimani

[View project](#)**Project**

A study of the rate and causes of addiction relapse among volunteer addicts seeking help at the Center for the Prevention of Addiction affiliated to the Welfare Organization, Ardebil province, Iran.

Mohammad Narimani · Abass Abolghasemi · الیه علمداری

1-how much is the rate of addiction relapse amoneg addicts self-introduced to the welfare organization of ardabil province ? 2-which social ,familial, personal factors are related to addiction rel ... [\[more\]](#)[View project](#)**Project**

cognitive fusion

Deleted Profile · Mohammad Narimani

[View project](#)**Article**

The prevalence of post traumatic stress disorder is increased among intensive care unit nurses

January 2006

Meredith Mealer · A.L. Shelton · Greg S Martin · [...] · Marc Moss

[Read more](#)**Article** [Full-text available](#)

Psychosocial recovery after serious injury

December 2014 · European Journal of Psychotraumatology

Meaghan O'Donnell

The 2010 iteration of the Global Burden of Disease statistics (Murray et al., 2012) points to the growing impact of injury and highlights the mounting burden of psychiatric disorder. It is essential to examine the intersection between these two contributors to disease burden. The Australian Injury Vulnerability Study collected data of over 1,000 injury patients from their initial hospitalization ... [\[Show full abstract\]](#)

[View full-text](#)[Article](#)**Company** Disorders of extreme stress following war-zone military trauma: Associated features of posttraumatic...[Support](#)[Business solutions](#)

February 1999 · Journal of Consulting and Clinical Psychology

[About us](#)[Help Center](#)[Advertising](#)[News](#)

JD Ford

[Recruiting](#)[Careers](#)

Disorders of extreme stress not otherwise specified (DESNOS) and posttraumatic stress disorder (PTSD) were found to be comorbid but distinct among military veterans seeking inpatient PTSD treatment: 31% qualified for both conditions, 29% were diagnosed PTSD only, 26% were classified DESNOS only, and 13% met criteria for neither. PTSD diagnosis was associated with elevated levels of war-zone ... [\[Show full abstract\]](#)

© 2008-2023 ResearchGate GmbH. All rights reserved.

[Terms](#) · [Privacy](#) · [Copyright](#) · [Imprint](#)[Article](#) [Full-text available](#)

Psychology of trauma: Vulnerability and resilience

January 2013

Elizabeth Batista Pinto Wiese

The exposures to traumatic experiences often have a massive interference in the individual's psychological development and mental health. These experiences must be considered within a culture, as the cultural context shapes their effects on life , attributing to them different values and characteristics including the ones that are considered traumatic. Therefore, also the models for trauma ... [\[Show full abstract\]](#)

[View full-text](#)