

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/309764613>

Effect of magnesium on growth, fruit quality and sugar content in cucumber under various light intensities

Article · September 2015

CITATION

1

READS

416

3 authors:



Rasoul Azarmi

University of Mohaghegh Ardabili

16 PUBLICATIONS 448 CITATIONS

SEE PROFILE



Seyed Jalal Tabatabaei

Shahed University

92 PUBLICATIONS 1,120 CITATIONS

SEE PROFILE



Nader Chaparzadeh

Azərbaycan Şahid Mədani University

65 PUBLICATIONS 930 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



"split root" View project



محللول پاشی View project



**EFFECT OF MAGNESIUM ON GROWTH, FRUIT QUALITY AND SUGAR
CONTENT IN CUCUMBER UNDER VARIOUS LIGHT INTENSITIES**

RASOUL AZARMI^{1*}, SEYED JALAL TABATABAEI², NADER CHAPARZADEH³

¹Department of Horticultural Sciences, Faculty of Agriculture, Tabriz University, Tabriz,
Iran

²Faculty of Agriculture, Shahed University, Tehran, Iran

³Department of Biology, Faculty of Science, University of Azarbaijan Shahid Madani,
Tabriz, Iran

***Corresponding Author: E Mail: r_azarmi@uma.ac.ir; Fax: +98 45 32715417**

ABSTRACT

The effect of various Mg concentrations (0, 1, 2, 3 and 4 mM) in the nutrient solution on plant growth, fruit quality and sugars content in hydroponically grown cucumber (*Cucumis sativus* cv. Nagen 792) under optimum (100%) and low (50%) light intensities was evaluated. The results showed that the decreases in the dry matter, SPAD index and soluble protein and accumulation of soluble sugar and starch content in the leaves Mg deficiency (0 mM Mg) are suggestive of decreased growth, and the decrease induced by Mg deficiency was bigger under low light intensity than under optimum light intensity. Plant growth was improved at 3 mM Mg, but it was reduced when the Mg concentration increased (4 mM Mg). Concentration of Mg in the leaf and fruit increased drastically with increasing Mg in the nutrient solution. This became steadily more pronounced under low light intensity. Mg deficiency plants (0 mM Mg) developed visible symptoms- interveinal necrosis in middle leaves, especially optimum light intensity. Fruit quality traits such as fruit dry matter percentage and total soluble solid (TSS) and fruit Fv/Fm were increased at higher concentration of Mg (3 mM) in the solution, especially under optimum light intensity. But, fruit firmness was improved at lower concentration of Mg (1 mM) in the solution especially in optimum light intensity. In conclusion, Mg requirement of cucumber plants likely increases with light intensity. Thus, higher concentration of Mg (3 mM) in the

nutrient solution was the most favorable for cucumber plant growth and fruit quality grown in hydroponics.

Keywords: Mg, Growth, Quality, Cucumber, Light intensity

INTRODUCTION

Mg is an essential element for plant growth and development and fruit quality. Apart from being a central atom of the chlorophyll molecule, Mg also acts as activator or regulator of many key enzymes in plant physiological processes [38, 50]. Both Mg deficiency and oversupply have detrimental effects on plant photosynthesis [49], consequently resulting in abnormal or restricted growth of plants [50]). Mg plays a fundamental roles in phloem export of photosynthates so that a deficiency of Mg restricts the partitioning of dry matter between roots and shoots to result in excessive sugar, starch and amino acid accumulation in leaves (source tissues), chlorophyll breakdown, an over- reduction in the photosynthetic electron transport chain and the generation of highly reactive oxygen species (ROS) because of impairment in photosynthetic CO₂ fixation [9,25].

Mg had a greater effect on quality parameters, when Mg was supplies to low-Mg plants. However, there were no additional benefits when Mg was supplied to plants already grown under adequate Mg supply conditions [8]. Increasing Mg

supply on Mg-deficiency plants caused to increase fruit total soluble solid (TSS), dry matter and juice acidity, [21]. Excessive supply of Mg to fruit has negative effects on fruit firmness, texture and storability that are mainly determined by its antagonistic relationship with Ca[36]. Despite the well-known fundamental roles of Mg in plant metabolism, there is very limited information on interactions between Mg and light intensity on fruit quality of cucumber.

The responses of plants to different Mg concentrations are not only affected by Mg availability in the root zone, but also depend on light intensity, temperature and species[10, 27]. The roles of Mg in plant metabolism particularly under stress conditions are well known [9]. The authors indicated that the Mg requirement is increased under high-light conditions. The higher requirement of Mg under high light might be reduced to the fact that under suboptimal Mg supply and high light processes are induced which finally lead to accumulation of reactive oxygen species (ROS) and thus plant damage. As plants are subjected to various light

intensities at different seasons, this may alter the ability of plants to take up and translocate Mg. It seems that the adjustment of Mg concentration in the nutrient solution according to the light intensity should be crucial. The objective of this experiment was to determine the effects of Mg and light intensity on cucumber growth and quality traits in hydroponics.

MATERIALS AND METHODS

Plant materials and growth conditions

The experiment was carried out at the Department of Horticultural Science, University of Tabriz, Iran. Cucumber (*Cucumis sativus* L. cv. Nagen 792) seeds were sown in cells plug trays filled with vermiculite, after emergence of two true leaves, seedlings were transplanted to a 14l growth bags (70, 20, 10 cm) filled with a mixture of perlite and vermiculite (1:1 v/v). The nutrient solution was prepared based on full strength of Hoagland's solution [26] containing: 5.6 mM Ca (NO₃)₂, 4 mM KNO₃, 1 mM KH₂PO₄. The solution pH was maintained close to 6.5 by adding H₂SO₄. The electrical conductivity (EC) of the nutrient solution was within the range 2.2 - 2.4 dS m⁻¹. In order to keep the anionic-cationic balance and a similar electrical conductivity for the five solutions, mineral concentrations were adjusted leading to only slight

variations. The greenhouse was under natural photoperiod condition during spring and summer and air temperature was set to 27 ± 2 °C and 18 ± 2 °C in the day and at night, respectively. The experiment was a split-plot design with light intensity as the main plot and various Mg concentrations as subplot with three replications in each treatment. Each plot contained three plants. The plants were treated with five Mg concentrations (0, 1, 2, 3 and 4 mM) as MgSO₄.7H₂O. Treatments were labelled Mg 0, Mg 1, Mg 2, Mg 3 and Mg 4. The plants were subjected to two light intensity treatments [optimum (100%) low (50%) light intensities] using green shade netting suspended above the box frame with the size of 1.5 m × 8 m × 4 m. The box frames were randomly placed in the greenhouse. Everyday light intensity at the canopy height under the shaded netting and in the glasshouse was monitored using a light-meter (Skye Instrument. Powys. UK). The average of light intensity under shaded netting and in the glasshouse (unshaded) over entire period of experimentation is shown in Fig.1.

Data collection and chemical analysis

At the end of the experiment, two plants from each treatment harvested and the internode diameter were recorded. The plant organs divided into leaf and stem,

weighed and then all plant parts dried at 80 °C in an air-forced oven for 48 h for determination of leaf and stem dry matter. The percentage leaf and stem dry weight was then calculated as below: [(Dry weight/ Fresh weight) ×100]. Chlorophyll index value of fully expanded young leaves was determined using a portable SPAD-502 meter (Minolta, Tokyo, Japan) during the period of the plant's growth. Fruit quality was measured in a representative sample collected at the same position from plants in each treatment. The samples were taken from fruits with the same size. Each fruit was cut in to pieces and homogenized in a conventional blender in order to obtain the fruit juice. Thereafter, the fruit juice was filtered using a Whitman No. 4 filter paper and the filtrate was used to determine the pH, EC and TSS. The TSS content of the fruit was determined by using a digital refractometer (Atago Co., Tokyo, Japan). The juice pH and EC was measured by pH meter and EC meter, respectively. The measurement fruit firmness was determined using a penetrometer (Model: ST 977. Italy). A thin layer of the middle of fruit skin (0.5 mm) was removed by a sharp razor and fruit color or the extent of greening was measured using a chlorophyll-meter (SPAD-502, Konica, Minolta, Osaka,

Japan). Titratable acidity was measured by titrating with 0.1 M NaOH to the neutralization point. Before chlorophyll fluorescence measurements, fruits were dark-adapted for 30 min using a dark towel. Measurements were taken in the middle and near the neck position positions of each fruit at the same location in the fruit surface and then averaged. The maximal quantum yield of PS II photochemistry (Fv/Fm) was measured using a plant efficiency analyzer, Handy PEA (Hansatech Instruments). Ten fruits per treatment per replicate were used for the determination of fruit dry weight. Each individual fruit from each treatment was placed in a sampling bag and dried in the oven at a temperature of 80°C for 48 h until a constant weight was obtained. The percentage dry weight was calculated as below: [(Dry weight/ Fresh weight) ×100].

Soluble sugars were extracted using the method described by Sheligl(1986). About 0.5 g of dried leaf samples were extracted three times in 5ml of hot 80 % ethanol (80 °C)[51]. The supernatants from each extraction were combined and made to a convenient volume. 1 ml 5 % (w/v) phenol and 5ml concentrated H₂SO₄ were added to 2 ml the plant extract and mixed thoroughly. The reaction mixture was allowed to stand for 30 min before

the absorbance was recorded at 485 using a spectrophotometer (Motic, CL-45240-00, Hong Kong, china). Total sugar content of the sample was calculated based on calibration curve from a glucose working standard. Starch content was extracted from the residual plant material from the soluble sugar extraction described above. This was done by incubating the dry pellet with 2 ml HCl (4.68M) in boiling water bath for 15 min. the soluble products were assayed by the same phenol-sulphuric method described above. Soluble protein content was determined in according to Bradford (1976) using bovine serum albumin as standard[5]. To measure the Mg, Leaves and fruits washed with distilled water were oven-dried at 80 °C for 48 h and weighed. The dry samples were ground to pass through a 0.5-mm screen. 1 g dry samples of leaves and fruits a were soaked in 10mL nitric acid (HNO₃) for 24 h then digested in digestion systems in a fume hood, heated to 110 °C for 3 h. The extracted solution was transferred to 100 mL volumetric flasks, and then diluted to 100mL with deionized water for Mg assays [35]. The Mg concentration in the leaf and fruit were measured at a wavelength 285.2 nm by atomic absorption spectrophotometry (Perkin-Elmer, Model 110, and USA).

Statistical Analysis

A statistical analysis was made using analysis of variance the SPSS 21 software and the means were separated by the Duncan test at a significance level of 0.05. The graphs were drawn using Excel software.

RESULTS AND DISCUSSION

Vegetative growth

The results showed that the highest leaf dry matter percentage was obtained in 3 and 4 mM Mg treatments. The leaf dry matter percent in concentration of 0 mM Mg was slightly higher than concentration of 1 mM Mg. Low light intensity largely decreased leaf dry matter percent compared to optimum light intensity (Table 1). This observation is in agreement with the finding of Lasa et al (2000) who showed that concentration of 0 mM Mg decreased 40 – 50% of shoot biomass compared with Mg sufficient plants in sunflower plants[32]. Low light intensity reduces the export of photosynthates from vegetative organs to the fruits. The reason for increase in leaf dry matter at 0 mM Mg may be due to impaired export of carbohydrates from source to sink sites and accumulation of soluble sugars in source leaves. Stem dry matter was not affected by various Mg concentrations, but optimum light intensity significantly increased stem dry

matter than low light intensity (Table 1). In the present experiment, there was a significant difference in leaf SPAD value between low Mg concentrations and sufficient Mg concentrations (Table 1). Significant decrease in chlorophyll concentration in Mg deficiency leaves has been widely reported [23, 55]. A reason for higher chlorophyll content under adequate Mg supply could be an enhanced production of chlorophyll and chlorophyll associated proteins. It is well documented that chlorotic and necrotic symptoms appearance in Mg deficiency leaves is associated with chlorophyll destruction due to photo-oxidation and accumulation of soluble sugar and starch in source leaves [7]. In both light intensities, internode diameter was increased with increasing Mg concentration in the solution. However, internode diameter was higher under optimum light intensity compared to low light intensity (Table 1). Light quantity and quality are major determinants of internode growth. Reductions in both photosynthetically active radiation (PAR), and red: far-red ratio (R: FR) result in similar shade avoidance responses, such as increased internode elongation and thicker internode. In plant communities, the R: FR ratio seems to act as an early competition signal [20, 31].

Soluble and insoluble sugars and soluble protein

The results showed that under both light intensities, concentration of 0 mM Mg in the solution had higher leaf soluble sugar and starch content compared to other treatments. However, leaf soluble sugar and starch content in optimum light intensity was higher than in low light intensity (Table 1). In almost all higher plants, the principle end products of leaf photosynthates are sucrose and starch. However, partitioning of sucrose and starch and their effect on dry matter distribution is influenced by several environmental factors, such as low temperature, drought and essential mineral nutrients [28, 56]. Mineral nutritional status of plants has a considerable impact on partitioning of carbohydrates and dry matter between shoots and roots [15, 34, 38]. Under Mg deficiency, starch concentrations are high in source leaves [18] and low in sink organs such as cereal grains and fruits [2]. This may demonstrate impaired photosynthate transport from source leaves to sink organs. Hence, in Mg-deficient plants higher shoot/root ratios were found compared with Mg-sufficient plants [4, 10, 16]. The translocation of amino acids and sugars from sink to source might be inhibited under magnesium deficiency

because of the effect of Mg on the H⁺-ATPase [9]. The results clearly showed that by increasing Mg concentration in the solution, increased soluble protein content in the cucumber leaves (Table 3). Andrews et al (1999) reported that Mg deficiency induced an increase in the protein content of *Pisum sativum* and *Phaseolus vulgaris*[1]. The reduction of protein in Mg deficiency plants could be attributed to a decrease in protein synthesis due to the participation of Mg in the aggregation of ribosome subunits and its requirement for RNA polymerases [12]. Protein biosynthesis also is strongly reduced under Mg deficiency leading to increased concentrations of the precursor amino acids [19,40].

Leaf and fruit Mg concentration

The results indicated that with increased Mg concentration in the nutrient solution led to a significant increase in Mg content in leaves and fruits under both light intensity. But, Mg concentration in cucumber leaves and fruits under low light intensity was higher than in cucumber leaves under optimum light intensity (Fig. 2). Greater concentration of Mg was observed with increasing Mg levels in the nutrient solution. However, Mg concentration of leaf in shaded plant was higher than in unshaded plants. Visual symptoms of Mg deficiency

appeared only in 0 mM Mg concentration and under both light intensities. However the symptoms severity became more pronounced under optimum light intensity. These symptoms observed after 35 days of treatment initiation and in middle leaves as necrotic lesion. Whereas, no visual symptoms of Mg deficiency were found in leaves of any other treatments in the range of 1 to 4 mM Mg concentrations, under both light intensity. The incidence of Mg deficiency was attenuated by the initial amount Mg present within the plant. Because the cucumber seedlings had been grown in one third of full nutrient solution (containing 0.3 mM Mg) for four weeks prior to treatment initiation, the initial accumulated Mg and its internal recycling in the seedling attenuated the visible signs of Mg deficiency. The Mg concentration sufficient for optimal growth varied with species. Kirkby and Mengel, (1979) reported that 0.35- 0.8% in the dry weight to be sufficient for cucumber[29]. However, the results obtained in this study agree well with the general threshold line for the occurrence of Mg deficiency determined by Kirkby and Mengel, (1979)[29]. The Mg- deficiency visible symptoms observed partially only on the full developed middle leaves [6, 7, 19, 43]. In cucumber, deficiency visible

symptoms observed initially as interveinal chlorosis and finally, as interveinal necrosis on leaves. The occurrence of Mg- deficiency on the middle leaves could significantly affect the photoassimilate production and supply to other parts of plants. Both shading and Mg levels in the nutrient solution altered Mg concentration in the leaves. This is consistent with findings by Zhao and Oosterhuis (1998) and Sonneveld (1987) who indicated that high light intensity will decrease the ability of plants to absorb and translocate magnesium, since transpiration is reduced and the translocation of magnesium is driven by transpiration rates [57, 54]. In general, the breakdown of chlorophyll under magnesium deficiency is associated with the accumulation of sugars and starch in the cells of deficient leaves [24, 9]. This causes an over-reduction of the photosynthetic electron transport chain, which leads to the formation of reactive oxygen species.

Fruit quality traits

Despite the well-known roles of Mg in plant metabolism, very limited information there have been concerning the significance of Mg for the quality of agriculture and horticulture produce, as compared to other major nutrients. A fruit quality trait like fruit firmness was significantly affected by treatments. So

that the firmest fruit under optimum light intensity in 1 and 2 mM Mg and in low light intensity at 1 mM Mg were obtained. Under optimum light intensity conditions, the fruits were firmer than under low light intensity conditions. This is consistent with findings by Marcelle, (1995) who showed that an optimal Mg concentration 'has to be relatively low' for good storage properties [38]. The reduced firmness in plant grown with low Mg content may be due to high concentration of fruit Ca. Fruit Firmness is an extremely important quality attribute of cucumber and consumer prefer a firm and crisp product. The Mg: Ca ratio mainly determines service and stability aspects as components of the total food quality, such as product firmness, texture and storability that are mainly determined by the role of Ca in stabilizing cell walls [21]. Since Mg is capable of replacing Ca from binding sites, imbalance Mg: Ca ratios in the tissue often negatively affect product quality. The fruit TSS increased with increasing Mg concentration up to 3 mM Mg and then decreased with increasing Mg (4 mM). The fruit TSS under optimum light intensity was higher than under low light intensity. The increase of fruit juice TSS with increasing Mg concentration in the solution reported by Quaggio et al (1992)

who indicated that juice pH, Soluble Solid and titratable acidity of fruit orange increased with increasing Mg[45]. Mg:K ratio appears to influence primarily quality properties through the role of both mobile cations in metabolite formation and translocation to fruits [46]. The highest marketable, high quality yield was observed when K and Mg were supplied in the highest amounts at a ratio of 5:1. This clearly points to the facts that only balanced crop nutrition can result in optimal quality [3]. Fruit juice pH and EC were not affected by the treatments (Table 3). Fruit dry matter increased with increasing Mg concentration in the solution. Low light intensity decreased fruit dry matter (Table 2). Dry matter content is an important quality criterion. Marcelis (1993) showed a positive relationship between the dry matter distribution of the fruit and irradiance[37]. Increases of dry matter percent can be due to the importance of Mg for photosynthesis and assimilate translocation. This finding was underlined by Feltran et al (2004) and Poberezny and Wszelaczynska (2001) who showed that increasing Mg supply consistently increased dry matter in potato[17, 44]. Also, Hao and Papadopoulos(2004) indicated that at a given Ca supply increasing the Mg

application enhanced the biomass allocation to the fruit, whereas the allocation to the leaves decreased, pointing to the decisive role of Mg in carbohydrate partitioning[22]. The highest value in fruit Fv/Fm was obtained in 2 and 3 mM Mg treatments. Fruit Fv/Fm was higher under low light intensity than that under optimum light intensity (Table 2). Chlorophyll fluorescence is an indirect measurement of the physiological status of green tissues [48, 41], being used in both green leaves and several chlorophyll-containing fruits [52, 53, 13]. Concerning the evaluation of changes in fruit tissues, chlorophyll fluorescence measurement has the advantage of detecting cellular injury due to natural senescence or environmental stresses in advance to the development of visible symptoms [14, 52, 47]. Fruit chlorophyll was significantly higher under optimum light intensity than under low light intensity (Table 2). Fruit color is one of the few practical criteria for assessing cucumber quality after harvest at present. A dark green cucumber is expected to have a longer shelf-life than a light green cucumber [33]. This result is agree with observation of Klieber et al (1993) who showed that high light intensity is necessary for high chlorophyll content in cucumber[30]. They also confirmed that high chlorophyll content

was positively correlated with a high percentage of PPF reaching the fruit surface.

CONCLUSION

It can be concluded that Mg requirement of cucumber plants likely increases with light intensity. The moderate concentration of Mg (2 mM) in the solution was the most desired for cucumber fruit quality. While, higher concentration of Mg (3 mM) in the solution was the most favorite for cucumber growth in hydroponics system.

REFERENCES

- [1] Andrews M, Sprent JI, Raven J A, Eady P E. (1999). Relationships between shoot to root ratio, growth and leaf soluble protein concentration of *Pisum sativum*, *Phaseolus vulgaris* and *Triticum aestivum* under different nutrient deficiencies. *Plant Cell and Environment* 22, 949–958.
- [2] Beringer H, Forster H. 1981. Einfluss variierter Mg-ernahrung auf tausendkorngewicht und P-fraktionen des gerstenkorns. *Z Pflanzenernahr Boden* 144: 8–15.
- [3] Borkowski J, Szwonek E. 1986. Effect of potassium and magnesium on the quality of tomato fruits. *Acta Horticulture* 191: 133–139.
- [4] Bouma D, Dowling EJ, Wahjoedi H. 1979. Some effects of potassium and magnesium on the growth of subterranean clover (*Trifolium subterraneum*). *Annals Botany* 43: 529–538.
- [5] Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye-binding. *Analytical Biochemistry* 72, 248–254.
- [6] Broschat TK. 1997. Nutrient distribution, dynamics, and sampling in coconut and Canary Island date plants. *Journal of the American Society for Horticultural Science* 122, 884–890.
- [7] Cakmak I. 1994. Activity of ascorbate-dependent H₂O₂-scavenging enzymes and leaf chlorosis are enhanced in magnesium- and Potassium-deficient leaves, but not in phosphorus-deficient leaves. *Journal of Experimental Botany* 278, 1259–1266.
- [8] Cakmak I. 2013. Magnesium in crop production, food quality and human health. *Plant Soil* 368: 1–4.
- [9] Cakmak I, Kirkby EA. 2008. Role of magnesium in carbon

- partitioning and alleviating photooxidative damage. *Physiologia Plantarum* 133: 692–704.
- [10] Cakmak I, Marschner H. 1992. Magnesium- deficiency and high light intensity enhance activities of superoxide dismutase, ascorbate peroxidase and glutathione reductase in bean leaves. *Plant Physiology* 98: 1222–1227.
- [11] Cakmak I, Yazici AM. 2010. Magnesium: a forgotten element in crop production. *Better Crops* 94: 23-25.
- [12] Cammarano P, Felsani A, Gentile M, Gualerzi C, Romeo C, Wolf G. 1972. Formation of active hybrid 80-S particles from subunits of pea seedlings and mammalian liver ribosomes. *Biochemistry and Biophysica Acta* 281: 625–642.
- [13] DeEll JR, Toivonen PMA. 2000. Chlorophyll fluorescence as a nondestructive indicator of broccoli quality during storage in modified-atmosphere packaging. *Hort Science* 35: 256–259.
- [14] DeEll JR, Prange RK, Murr DP. 1997. Chlorophyll fluorescence as an indicator of apple fruit firmness. In: *Proceedings of the International Conference on Sensors for Nondestructive Testing: Measuring the Quality of Fresh Fruits and Vegetables*, p. 67–73.
- [15] Druege U, Zerche S, Kadner R, Ernst M. 2000. Relation between nitrogen status, carbohydrate distribution and subsequent rooting of chrysanthemum cuttings as affected by pre-harvest nitrogen supply and cold-storage. *Annals of Botany* 85(5): 687-701.
- [16] Ericsson T. 1995. Growth and shoot root ratio of seedlings in relation to nutrient availability. *Plant and Soil* 168/169: 205-14.
- [17] Feltran JC, Lemos LB, Vieites RL. 2004. Technological quality and utilization of potato tubers. *Scientia Agricola* 61: 593–597.
- [18] Fischer ES, Bremer E. 1993. Influence of magnesium deficiency on rates of leaf expansion, starch and sucrose accumulation, and net assimilation in *Phaseolus vulgaris*. *Physiologia Plantarum* 89: 271-276.
- [19] Fischer ES, Lohaus G, Heineke D, Heldt HW. 1998. Magnesium

- deficiency results in accumulation of carbohydrates and amino acids in source and sink leaves of spinach. *Physiologia Plantarum* 102: 16-20.
- [20] Franklin KA, Whitlam GC. 2005. Phytochromes and shade-avoidance responses in plants. *Annals of Botany* 96: 169–175.
- [21] Gerendas J, Fuhrs H. 2013. The significance of magnesium for crop quality. *Plant Soil* 368: 101-128.
- [22] Hao X, Papadopoulos AP. 1999. Effects of supplemental lighting and cover materials on growth, photosynthesis, biomass partitioning, early yield and quality of greenhouse cucumber. *Scientia Horticulture* 80: 1-18.
- [23] Hariadi Y, Shabala S. 2004. Screening broad beans (*Vicia faba*) for magnesium deficiency. II. Photosynthetic performance and leaf bioelectrical responses. *Functional Plant Biology* 31: 539-549.
- [24] Hermans C, Verbruggen N. 2005. Physiological characterization of Mg deficiency in *Arabidopsis thaliana*. *Journal of Experimental Botany* 56: 2153-2161.
- [25] Hermans C, Bourgis F, Faucher M, Delrot S, Strasser RJ, Verbruggen N. 2005. Magnesium deficiency in sugar beet alters sugar partitioning and phloem loading in young mature leaves. *Planta* 220: 441-449.
- [26] Hoagland DR, Arnon DS. 1950. The water culture method for growing plants without soil. California Agricultural Experiment Station Circular 374: 1–32.
- [27] Huang JW, Welch RM, Grunes DL. 1990. Magnesium, nitrogen form, and root temperature effects on grass tetany potential of wheat forage. *Agronomy Journal* 82(3): 581–587.
- [28] Huber SC. 1989. Biochemical Mechanism for Regulation of Sucrose Accumulation in Leaves during Photosynthesis. *Plant physiology* 91(2): 656- 662.
- [29] Kirkby EA, Mengel K. 1979. The role of magnesium in plant nutrition. - *Z. Pflanzenemahr. Bodenkd* 2: 209-222
- [30] Klieber A, Lin WC, Joliffe PA, Hall JW. 1993. Training methods affect canopy light exposure and shelf life of long English cucumber. *Journal of American*

- Society Horticulture Science 118, 786–790.
- [31] Kurepin LV, Emery RJN, Pharis RP, Reid DM. 2007. Uncoupling light quality from light irradiance effects in *Helianthus annuus* shoots: putative roles for plant hormones in leaf and internode growth. *Journal of Experimental Botany* 58: 2145–2157.
- [32] Lasa B, Frechilla S, Aleu M, Gonzales-Moro B, Lamsfus C, Aparicio-Tejo PM. 2000. Effects of low and high levels of magnesium on the response of sunflower plants grown with ammonium and nitrate. *Plant and soil* 225:167-174.
- [33] Lin WC, Ehret DL. 1991. Nutrient concentration and fruit thinning affect shelf life of long English cucumber. *Hortscience* 26 (10): 1299–1300.
- [34] López-Bucio J, Cruz-Ramirez A, Herrera-Estrella L. 2003. The role of nutrient availability in regulating root architecture. *Current Opinion in Plant Biology* 6(3): 280-287.
- [35] MAFF. 1985. The analysis of agricultural materials. Ministry of Agriculture Fisheries and Food, London, UK.
- [36] Malundo TMM, Shewfelt RL, Scott JW. 1995. Flavor quality of fresh tomato (*Lycopersicon esculentum* Mill.) as affected by sugar and acid levels. *Post Biology Technology* 6:103–110.
- [37] Marcelis LFM. 1993. Fruit growth and biomass allocation to the fruits in cucumber. 1. Effect of fruit load and temperature. *Scientia Horticulturae* 54: 107–121.
- [38] Marcelle RD. 1995. Mineral nutrition and fruit quality. *Acta Horticulture* 383:219–226.
- [39] Marschner H. 1995. Mineral nutrition of higher plants. 2nd end. (Academic Press. San Diego, CA)
- [40] Marschner P. (ed) 2012. Marschner mineral nutrition of higher plants (Third Edition). Elsevier Ltd.
- [41] Maxwell K, Johnson GN. 2000. Chlorophyll fluorescence a practical guide. *Journal of Experimental Botany* 51: 659–668.
- [42] Mengel K, Kirkby EA. 1987. Principles of plant nutrition, 4th Edition, International Potash Institute, Switzerland.

- [43] Papenbrock J, Pfundel E, Mock HP, Grimm B. 2000. Decreased and increased expression of the subunit CHL1 diminishes Mg chelatase activity and reduces chlorophyll synthesis in transgenic tobacco plants. *The Plant Journal* 22: 155-164
- [44] Poberezný J, Wszelaczyńska E. 2011. Effect of bioelements (N, K, Mg) and long-term storage of potato tubers on quantitative and qualitative losses Part II. Content of dry matter and starch. *Journal of Elementology* 16: 237–246.
- [45] Quaggio JA, Sobrinho JT, Dechen AR. 1992. Magnesium influences on fruit yield and quality of ‘Valencia’ sweet orange on Rangpur lime. *Proceedings International Society Citriculture* 2: 633–637.
- [46] Romheld V, Kirkby EA. 2007. Magnesium functions in crop nutrition and yield. *Proceedings of a Conference in Cambridge* (7th Dec. 2007), 151–171.
- [47] Sanxter SS, Yamamoto HY, Fisher DG, Chan HT. 1992. Development and decline of chloroplasts in exocarp of *Caricacpapaya*. *Canadian Journal Botany* 70: 364–373.
- [48] Schreiber U, Bilger W. 1987. Rapid assessment of stress effects on plant leaves by chlorophyll fluorescence measurements. In: Tenhunen, J.D., Catarino, F.M., Lange, O.L., Oechel, W.C. (Eds.), *Plant response to stress-functional analysis in Mediterranean ecosystems*. NATO advanced science institute Series. Springer-Verlag Berlin Heidelberg New York p. 27–53.
- [49] Shabala S, Hariadi Y. 2005. Effects of magnesium availability on the activity of plasma membrane ion transporters and light-induced responses from broad bean leaf mesophyll. *Planta* 221(1): 56-65.
- [50] Shaul O. 2002. Magnesium transport and function in plants; the tip of the iceberg. *Biology of Metals* 15: 309- 323.
- [51] Sheligl HQ. 1986. Die verwertung orgngischer souren durch chlorella lincht. *Planta Journal* 47-51.
- [52] Smillie RM, Hetherington SE, Nott R, Chaplin GR, Wade NL. 1987. Application of chlorophyll

fluorescence to the postharvest physiology and storage of mango and banana fruit and the chilling tolerance of mango cultivars. *Asian Food Journal* 39: 55–59.

[53] Song F, Deng W, Beaudry RM. 1997. Changes in chlorophyll fluorescence of apple fruit during maturation, ripening, and senescence. *HortScience* 32: 891–896.

[54] Sonneveld C. 1987. Magnesium deficiency in rockwool-grown tomatoes as affected by climatic conditions and plant nutrition. *Journal of Plant Nutrition* 10: 1591–1604.

[55] Teklic T, Vrataric M, Sudaric A, Kovacevic V, Vukadinovic V, Bertic B. 2009. Relationship among chloroplast pigments concentration and chlorophyll meter readings in soybean under influence of foliar magnesium application. *Communications in Soil Science and Plant Analysis* 40: 706-725.

[56] Wardlaw A. 1990. The control of carbon partitioning in plants. *New Phytologist* 116(3): 341-381.

[1] Zhao D, Oosterhuis DM. 1998. Influence of shade on mineral nutrient status of field-grown cotton. *Journal of Plant Nutrition* 21(8): 1681–1695,

Table 1: The effect of Mg and light intensity on cucumber vegetative growth

| Light intensity | Mg (mM) | Leaf Dwt (%) | Stem Dwt (%) | Internode diagonal (cm) | Leaf SPAD index | Soluble sugar (mg g ⁻¹ DW) |
|--------------------|---------|--------------|--------------|-------------------------|---------------------|---------------------------------------|
| Optimum Light | 0 | 14.16 | 7.60 | 3.85 | 55.30 ^{ef} | 35.82 |
| | 1 | 13.55 | 7.85 | 4.46 | 58.70 ^e | 31.41 |
| | 2 | 14.44 | 7.80 | 4.54 | 59.50 ^{bc} | 25.42 |
| | 3 | 16.63 | 8.02 | 4.79 | 61.30 ^{ab} | 23.73 |
| | 4 | 15.36 | 7.75 | 4.11 | 62.63 ^a | 24.39 |
| Low Light | 0 | 11.23 | 6.29 | 3.83 | 53.76 ^f | 28.18 |
| | 1 | 11.20 | 6.66 | 4.05 | 54.33 ^{ef} | 24.73 |
| | 2 | 11.74 | 7.00 | 4.22 | 55.06 ^{ef} | 20.35 |
| | 3 | 13.17 | 6.94 | 4.00 | 56.23 ^{de} | 19.48 |
| | 4 | 12.13 | 6.60 | 3.83 | 57.43 ^{cd} | 18.74 |
| Optimum Light | | 14.83 | 7.80 | 4.35 | 59.48 | 28.15 |
| Low Light | | 11.89 | 6.70 | 3.98 | 55.36 | 22.29 |
| Light intensity | | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| Mg | | 0.01 | ns | 0.01 | 0.001 | 0.001 |
| Light intensity×Mg | | ns | ns | ns | 0.05 | ns |

(ns) non significance; (0.05) significance at 0.05 probability level; (0.01) significance at 0.01 probability level; (0.001) significance at 0.001 probability level. Within each column, same letters indicate no significant difference between treatments (P < 0.01)

Table 2: The effect of Mg and light intensity on cucumber fruit quality traits

| Light intensity | Mg (mM) | Firmness (Kg) | Fruit dry matter (%) | Fruit Fv/Fm value | Fruit color (SPAD UNIT) |
|--------------------|---------|--------------------|----------------------|-------------------|-------------------------|
| Optimum Light | 0 | 3.66 ^{bc} | 3.92 | 0.737 | 53.83 |
| | 1 | 3.73 ^{ab} | 4.83 | 0.744 | 56.40 |
| | 2 | 4.20 ^a | 4.35 | 0.753 | 57.20 |
| | 3 | 3.30 ^{bc} | 4.90 | 0.760 | 56.96 |
| | 4 | 3.26 ^{bc} | 4.71 | 0.741 | 55.94 |
| Low Light | 0 | 3.10 ^{bc} | 4.21 | 0.749 | 51.10 |
| | 1 | 3.66 ^a | 4.31 | 0.751 | 53.96 |
| | 2 | 3.63 ^{bc} | 4.48 | 0.763 | 53.23 |
| | 3 | 3.23 ^{bc} | 4.46 | 0.764 | 53.03 |
| | 4 | 3.33 ^{bc} | 4.67 | 0.751 | 50.76 |
| Optimum Light | | 3.63 | 4.85 | 0.747 | 56.07 |
| Low Light | | 3.39 | 4.42 | 0.756 | 52.42 |
| Light intensity | | 0.01 | ns | 0.01 | 0.01 |
| Mg | | 0.001 | 0.001 | 0.01 | ns |
| Light intensity×Mg | | 0.05 | ns | ns | ns |

(ns) non significance; (0.05) significance at 0.05 probability level; (0.01) significance at 0.01 probability level; (0.001) significance at 0.001 probability level. Within each column, same letters indicate no significant difference between treatments ($P < 0.01$)

Table 3: The effect of Mg and light intensity on cucumber fruit quality traits

| Light intensity | Mg (mM) | Fruit juice pH | Fruit juice EC (dS m ⁻¹) | Acidity (%) | Soluble protein (mg g ⁻¹ FW) | Fruit Mg (mgg ⁻¹ DW) |
|--------------------|---------|----------------|--------------------------------------|-------------|---|---------------------------------|
| Optimum Light | 0 | 5.93 | 0.50 | 1.993 | 1.00 | 0.720 |
| | 1 | 5.86 | 0.56 | 2.213 | 1.22 | 1.253 |
| | 2 | 5.93 | 0.56 | 2.327 | 1.40 | 2.133 |
| | 3 | 5.74 | 0.56 | 2.417 | 1.57 | 2.727 |
| | 4 | 5.83 | 0.56 | 2.650 | 1.61 | 3.047 |
| Low Light | 0 | 5.72 | 0.46 | 1.703 | 0.83 | 0.880 |
| | 1 | 5.64 | 0.56 | 1.960 | 0.91 | 1.480 |
| | 2 | 5.69 | 0.56 | 2.027 | 1.24 | 2.687 |
| | 3 | 5.55 | 0.55 | 2.193 | 1.45 | 2.753 |
| | 4 | 5.63 | 0.56 | 2.387 | 1.43 | 3.213 |
| Optimum Light | | 5.86 | 0.55 | 2.320 | 1.36 | 1.97 |
| Low Light | | 5.65 | 0.54 | 2.054 | 1.17 | 2.16 |
| Light intensity | | 0.05 | ns | ns | 0.01 | 0.01 |
| Mg | | ns | ns | ns | 0.001 | 0.001 |
| Light intensity×Mg | | ns | ns | ns | ns | ns |

(ns) non significance; (0.05) significance at 0.05 probability level; (0.01) significance at 0.01 probability level; (0.001) significance at 0.001 probability level. Within each column, same letters indicate no significant difference between treatments ($P < 0.01$)




Fig. 1: Light intensity at the optimum (100%) and low (50%) light intensity during the entire of period of experimentation




Fig. 2: The effect of Mg and light intensity on the leaf Mg concentration in cucumber plants (error bars on the columns represent standard error)




Fig. 3: The effect of Mg and light intensity on fruit TSS in cucumber plants (error bars on the columns represent standard error)




Fig. 4: The effect of Mg and light intensity on starch content in cucumber plants (error bars on the columns represent standard error)




Fig. 5: Visual symptoms of Mg sufficient leaves (A) and Mg deficiency leaf (B)